

## Punto

**Seed treatment insecticide for use on cereals, forage brassicas, grass seed, maize, sweetcorn, potatoes, pumpkins and winter squash.**

**ACTIVE INGREDIENT:** Contains 600 g/L imidacloprid in the form of a suspension concentrate.

- Reduces the amount of foliar insecticides required to control the same target pests and saves resources because the crop does not need to be monitored as often
- Punto provides increased sustainability over granular and foliar insecticides due to being applied on the seed only which reduces residuals
- Punto is systemically transported throughout the plant during the development stages to ensure long lasting protection for the plant and help reach maximum productivity



### **Withholding periods:**

- Cereals, forage brassicas, pasture, maize and sweetcorn – 42 days
- Grass seed oversown into grazing pasture (stitching) – 21 days from oversowing

### **Application and directions of use:**

Shake or roll the container thoroughly before use. When decanting ensure adequate agitation is maintained. At the time of treatment, the seeds must be dyed a distinctive colour.

Apply using well maintained and accurately calibrated seed treatment equipment. As Punto may alter the seed flow rate, calibrate the planter prior to planting with treated seed.

**Use situations:**

<b>Crop</b>	<b>Pest</b>	<b>Rate</b>	<b>Notes</b>
Cereals	Grass grub larvae Aphids	90ml per 100kg seed	Apply with sufficient water (usually 10 litres/tonne of seed) to evenly cover the seed.
Forage brassicas	Aphids Argentine stem weevil Springtails (Sminthuridae family) Wheat bug	12-24ml per kg of seed	Apply to seed ensuring even coverage. The high rate will protect against aphids for longer.
Grass sees	Grass grub larvae Black beetle adults	580ml per 100kg seed	Apply with sufficient water (usually 10 litres/tonne) to evenly cover the seed.
Maize Sweetcorn	Black beetle Argentine stem weevil	580ml per 100kg seed	Apply with sufficient water (usually 1-2 litres/100kg of seed) to evenly cover the seed. Temporary phytotoxic effects may occur on some sweetcorn varieties, especially under dry conditions. These will disappear as the plant grows.
Potatoes	Aphids	90ml per tonne of seed	Apply with sufficient water (usually 2-3 litres/tonne) to evenly cover seed tubers. If treated tubers are to be stored before planting ensure they are completely dry before storing.
Pumpkins Winter squash	Aphids Argentine stem weevil Grass grub beetle Lucerne flea Springtails	50ml per kg of seed	Apply with sufficient water (usually 1-2 litres/100kg of seed) to evenly cover the seed.

**Product Codes:**

<b>Code</b>	<b>Product</b>	<b>Unit</b>	<b>Qty per Unit</b>	<b>Barcode</b>
AGPU0005	PUNTO SC SEED TREATMENT INSECTICIDE - 5L	EACH	1	9400579558048

**Health and Safety Information:**

Material safety data sheets are available on request.

Refer to product label for complete usage, safety and product information before use.

Registered pursuant to the ACVM Act 1997, P8806